

§ 520.905b

21 CFR Ch. I (4–1–05 Edition)

the indications provided in paragraph (d) of this section and for treating infections of stomach bot as provided in § 520.2520, have been shown to be compatible and not to interfere with one another.

[42 FR 59069, Nov. 15, 1977; 43 FR 12311, Mar. 24, 1978. Redesignated at 44 FR 1375, Jan. 5, 1979, and amended at 46 FR 29464, June 2, 1981; 47 FR 15327, Apr. 9, 1982; 48 FR 42809, Sept. 20, 1983; 49 FR 1983, Jan. 17, 1984; 53 FR 40058, Oct. 13, 1988; 59 FR 26943, May 25, 1994; 61 FR 29478, June 11, 1996; 63 FR 63983, Nov. 18, 1998; 66 FR 47960, Sept. 17, 2001; 68 FR 26205, May 15, 2003]

§ 520.905b Fenbendazole granules.

(a) *Specifications.* Each gram of granules contains 222 milligrams (mg) fenbendazole.

(b) *Sponsor.* See No. 057926 in § 510.600(c) of this chapter.

(c) *Special considerations.* See § 500.25 of this chapter.

(d) *Conditions of use*—(1) *Horses*—(i) *Amount.* 5 mg/kilogram (kg) for large strongyles, small strongyles, and pinworms; 10 mg/kg for ascarids.

(ii) *Indications for use.* For the control of infections of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles, pinworms (*Oxyuris equi*), and ascarids (*Parascaris equorum*).

(iii) *Limitations.* Sprinkle the appropriate amount of drug on a small amount of the usual grain ration. Prepare for each horse individually. Withholding feed or water is not necessary. Retreat in 6 to 8 weeks if required. Do not use in horses intended for food.

(2) *Dogs*—(i) *Amount.* 50 mg/kg daily for 3 consecutive days.

(ii) *Indications for use.* For the removal of ascarids (*Toxocara canis*, *Toxascaris leonina*), hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*), whipworms (*Trichuris vulpis*), and tapeworms (*Taenia pisiformis*).

(iii) *Limitations.* Mix the appropriate amount of drug with a small amount of the usual food; dry dog food may require slight moistening to facilitate mixing. Medicated food must be fully consumed.

(3) *Zoo and wildlife animals*—(i) *Amount.* 10 mg/kg per day for 3 days.

(ii) *Indications for use.* For control of internal parasites of *Felidae* and *Ursidae* as follows:

(A) Lion (*Panthera leo*) and Tiger (*Panthera tigris*): Ascarid (*Toxocara cati*, *Toxascaris leonina*), Hookworm (*Ancylostoma* spp.).

(B) Cheetah (*Acinonyx jubatus*): Ascarid (*Toxocara cati*, *Toxascaris leonina*).

(C) Puma (*Felis concolor*), Panther (*Panthera* spp.), Leopard (*Panthera pardus*), Jaguar (*Panthera onca*): Ascarid (*Toxocara cati*, *Toxascaris leonina*), Hookworm (*Ancylostoma* spp.), Tapeworm (*Taenia hydatigena*, *T. krabbei*, *T. taeniaeformis*).

(D) Black Bear (*Ursus americanus*): Ascarid (*Baylisascaris transfuga*, *Toxascaris leonina*), Hookworm (*Ancylostoma caninum*), Tapeworm (*Taenia hydatigena*, *T. krabbei*).

(E) Polar Bear (*Ursus maritimus*) and Grizzly Bear (*Ursus horribilis*): Ascarid (*Baylisascaris transfuga*, *Toxascaris leonina*).

(iii) *Limitations.* Top dress or mix with a small portion of food. Must be fully consumed prior to feeding. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Do not use 14 days before or during the hunting season.

[44 FR 1375, Jan. 5, 1979, as amended at 47 FR 15327, Apr. 9, 1982; 48 FR 50528, Nov. 2, 1983; 59 FR 35252, July 11, 1994; 66 FR 47960, Sept. 17, 2001; 67 FR 47450, July 19, 2002]

§ 520.905c Fenbendazole paste.

(a) *Specifications.* The product is an aqueous paste containing 10 percent fenbendazole.

(b) *Sponsor.* See No. 057926 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Conditions of use*—(1) *Horses*—(i) (a) *Amount.* 2.3 milligrams per pound of body weight (one 2.5-gram fenbendazole syringe for a 1,100-pound horse). For foals and weanlings (less than 18 months of age), 4.6 milligrams per pound of body weight (one 2.5-gram fenbendazole syringe for each 550 pounds of body weight).

(b) *Indications for use.* For control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles, pinworms (*Oxyuris equi*),